

Subharmonic Solutions for Second Order Differential Systems

NORIMICHI HIRANO

*Department of Mathematics, Yokohama National University,
Yokohama 240 Hodogaya, Japan*

Submitted by Jack K. Hale

Received December 6, 1993

Ink this paper, we consider the existence of subharmonic solutions for the problem $u_{tt} + G'(u) = f(t)$, where $G: R^N \rightarrow R$ is not necessarily convex and $f: R \rightarrow R^N$ is periodic with minimal period $T > 0$. © 1995 Academic Press, Inc.

1. INTRODUCTION

In this paper, we consider the system of second order differential equations of the form

$$(P) \quad u_{tt} + G'(u) = f(t),$$

where $G \in C^2(R^N; R)$ and $f: R \rightarrow R^N$ is a continuous function with minimal period $T > 0$. Here G' denote the gradient of G .

Our purpose in this paper is to show the existence of subharmonic solutions of (P). That is, we seek for a sequence $\{u_{k_i}\}$ of solutions of (P) where u_{k_i} is a periodic function with minimal period $k_i T$ and $k_i \rightarrow \infty$, as $i \rightarrow \infty$.

The existence of subharmonic solutions of (P) has been studied by many authors [1–6, 9, 11, 13] (see [5] for further references). Most of the existence results for subharmonic solutions assume the (strict) convexity of functional G , and few seem to be known in the case that G is not a convex function. In [7], Giannoni considered problem (P) with $G(x)$ replaced by $G(t, x): R \times R^N \rightarrow R$ which is not necessarily convex for the second variable and $\lim_{|x| \rightarrow \infty} G(t, x) = 0$ uniformly in t . Recently, Fonda and Lazer [5] established an existence result without assuming convexity on G . They proved the existence of subharmonic solutions of (P) in the case that G' is bounded and $\lim_{|x| \rightarrow \infty} G(x) = \infty$. Their argument

is based on the estimate of critical levels of functionals associated to problem (P). Here we also consider the case that functional G is not necessarily convex, under different assumptions from those of Fonda & Lazer. Instead of making use of the estimate of critical levels of functionals associated to problem (P), we show the existence of subharmonic solutions by applying a Morse index theorem for critical points of min-max type established by Lazer and Solimini [8] (cf. [6] for applications of Lazer and Solimini's results to the convex case). Our approach enables us to treat the case that G is bounded and the case that G' is unbounded.

2. PRELIMINARY NOTATIONS AND THE RESULTS

For each $T > 0$, we set

$$H_T = H^{1/2}(S_T^1; R^N),$$

where $S_T^1 = [0, T]/\{0, T\}$. H_T is regarded as a subspace of $L^2([0, T]; R^N)$. $\|\cdot\|_T$ and $\langle \cdot, \cdot \rangle_T$ stand for the norm and inner product of $L^2([0, T]; R^N)$, respectively. For each functional $f \in C^1(H_T, R)$ and $u \in H_T$, we denote by $f'(u)(v)$ the value of the gradient $f'(u): H_T \rightarrow R$ at $v \in H_T$. Let $v \in H_T$. We denote by \bar{v} the mean value of v , i.e.,

$$\bar{v} = (1/T) \int_0^T v(t) dt.$$

We also put $\tilde{v} = v - \bar{v}$. We define subspaces \tilde{H}_T and \overline{H}_T by

$$\tilde{H}_{kT} = \{v \in H_{kT} : \bar{v} = 0\} \quad \text{and} \quad \overline{H}_{kT} = \{v \in H_{kT} : \tilde{v} = 0\},$$

respectively. For each $x \in R^N$ and each $n \times n$ matrix A , we denote by $|x|$ and $|A|$ the norm of x and the operator norm of A , respectively. The inner product of R^N is denoted by $\langle \cdot, \cdot \rangle$. For $G: R^N \rightarrow R$, G'' denote the Hessian of G .

We can now state our main results.

THEOREM 1. *Assume that each critical point of G is nondegenerate and the following conditions are satisfied:*

- (G1) $\lim_{|x| \rightarrow \infty} G(x)/|x|^2 = 0$ and $\lim_{|y| \rightarrow \infty} G(y) > G(x)$ for all $x \in R$;
 (G2) $\lim_{|x| \rightarrow \infty} |G''(x)| = \lim_{|x| \rightarrow \infty} |G''(x)|/|G'(x)| = 0$.

Then there exist $T_0 > 0$ and $\rho > 0$ such that for each $f \in \tilde{H}_T$ with $T < T_0$ and $\|\tilde{f}\|_T < \rho$, problem (P) has periodic solutions with minimal period kT , for any sufficiently large prime number k .

Remark 1. The mapping G' can be unbounded under the hypotheses of Theorem 1. For example, conditions (G1) and (G2) are satisfied if $G(x) = |x|^p$ (or $G(x) = \sum_{i=1}^N |x_i|^p$) ($0 < p < 2$) for $|x|$ sufficiently large. For each $f \in H_T$, problem (P) can be rewritten as

$$u_{tt} + G'(u) - \bar{f} = \tilde{f}.$$

Then the assertion of Theorem 1 holds for each $f \in H_T$ if the assumptions of Theorem 1 are satisfied with G replaced by $G(x) - \langle \tilde{f}, x \rangle$. The nondegeneracy of G and condition (G2) imply implicitly that $G'(x) \neq 0$ if $|x|$ is sufficiently large.

THEOREM 2. Assume that G satisfies (G1), (G2). In addition, assume that

(G3) there exists $x_0 \in R^N$ such that

$$\langle G''(y)x_0, x_0 \rangle > 0 \quad \text{for all } y \in R^N. \quad (2.1)$$

Then there exists $T_0 > 0$ such that for each $f \in \tilde{H}_T$ with $T < T_0$, problem (P) has periodic solutions with minimal period kT , for any sufficiently large prime number k .

Remark 2. Condition (G3) is a directional convexity of G . That is, for any $z \in R^N$, the functional G is strictly convex on the line $\{z + tx_0 : t \in R\}$.

We can remove the restriction on the period T of f if we impose the boundedness condition on G' instead of G'' .

THEOREM 3. Assume that (G1) and (G3) hold. In addition, assume that

$$(G2') \quad \sup_{x \in R^N} |G'(x)| < \infty \quad \text{and} \quad \lim_{|x| \rightarrow \infty} |G''(x)|/|G'(x)| = 0.$$

Then for each $f \in \tilde{H}_T$ with $T > 0$, problem (P) has periodic solutions with minimal period kT , for any sufficiently large prime number k .

In the following, we assume that G satisfies the assumptions of Theorem 1. For each $x \in R^N$, $\lambda_1(x) \leq \lambda_2(x) \leq \cdots \leq \lambda_N(x)$ denote the eigenvalues of the matrix $G''(x)$. We denote by $x_{(i)}$ the i th element of $x \in R^N$. $D(r)$ stands for an open ball in R^N centered at 0 with radius r . From (G1), we can see that there exists a critical point of G . Since each critical point of G is nondegenerate and the set of critical points is bounded by assumption (G2), we can see that the set of critical points of G consists of finite elements.

Let $\{z_1, \dots, z_m\} \subset R^N$ be the set of critical points of G . Then there exists $c_0 > 0$ such that

$$|\lambda_j(z_i)| > 2c_0 \quad \text{for } 1 \leq i \leq m \text{ and } 1 \leq j \leq N.$$

For each $1 \leq i \leq m$ and $1 \leq j \leq N$, we denote by $w_{i,j}$ a normalized eigenvector corresponding to the eigenvalue $\lambda_j(z_i)$ of $G''(z_i)$. Then it follows that

$$|\langle G''(z_i)w_{i,j}, w_{i,j} \rangle| \geq 2c_0 \quad \text{for } 1 \leq i \leq m \text{ and } 1 \leq j \leq N. \quad (2.2)$$

Let $x, y \in R^N$. Then we have

$$G'(y) - G'(x) = \int_0^{|y-x|} G'' \left(x + t \left(\frac{y-x}{|y-x|} \right) \right) \frac{y-x}{|y-x|} dt. \quad (2.3)$$

On the other hand, we have by condition (G2) that for any $\delta > 0$, there exists $d_0(\delta) > 0$ such that

$$|G''(z)| \leq \delta |G'(z)| \quad \text{for all } z \in R^N \text{ with } |z| \geq d_0(\delta). \quad (2.4)$$

Now let $\varepsilon > 0$ and let $d(\varepsilon) > 0$ be so large that $d(\varepsilon) > d_0(\delta) + \varepsilon$, where $\delta > 0$ satisfies $(e^{\delta\varepsilon} - 1) = 1/2N$. Let $x, y \in R^N$ with $|x| > d(\varepsilon)$ and $y \in x + D(\varepsilon)$. Then since the line segment connecting x and y is contained in the set $\{z \in R^N : |z| > d_0(\delta)\}$, we find by (2.3) and (2.4) that

$$|G'(y)| \leq |G'(x)| + \delta \int_0^{|y-x|} \left| G' \left(x + t \left(\frac{y-x}{|y-x|} \right) \right) \right| dt.$$

It then follows from Gronwall's inequality that

$$|G'(y)| \leq e^{\delta|y-x|} |G'(x)|. \quad (2.5)$$

Then by (2.3), (2.4), and (2.5),

$$|G'(y) - G'(x)| \leq \delta \int_0^{|y-x|} e^{\delta t} |G'(x)| dt \leq (e^{\delta\varepsilon} - 1) |G'(x)| = |G'(x)|/2N$$

for all $x \in R^N$ with $|x| > d(\varepsilon)$ and $y \in x + D(\varepsilon)$. On the other hand, we have that for each $x \in R^N$, there exists i such that $|G'(x)_{(i)}| \geq |G'(x)|/N$.

Then it follows from the observation above that for each $\varepsilon > 0$, there exists $i \in \{1, \dots, N\}$ such that

$$G'(x)_{(i)} G'(y)_{(i)} > 0 \quad \text{for all } x \in R^N \text{ with } |x| > d(\varepsilon) \text{ and } y \in x + D(\varepsilon). \quad (2.6)$$

On the other hand, it follows from (2.2) that there exists $\varepsilon_0 > 0$ such that for each $1 \leq i \leq m$,

$$|G''(x) - G''(z_i)| < c_0/2(N-1) \quad \text{and} \quad |\langle G''(x)w_{i,j}, w_{i,j} \rangle| \geq c_0 \quad (2.7)$$

for all $1 \leq j \leq N$ and $x \in z_i + D(\varepsilon_0)$. We set

$$m_0 = \inf\{|G'(x)| : x \in R^N, |x| \leq d(\varepsilon_0), x \notin \bigcup (z_i + D(\varepsilon_0/2))\}. \quad (2.8)$$

It is obvious from the assumption that $m_0 > 0$.

Let E be a Banach space and $E = Y \oplus Z$, where Y and Z are subspaces of E with $\dim Y = k$. Let $F: E \rightarrow R$ be a C^2 functional. Let $u \in E$ be a critical point of F . If there exists a maximal integer q such that $F''(u): E \rightarrow E$ is negative definite (semi-negative definite) on some q -dimensional subspace of E , then the Morse index (augmented Morse index) of u is q . In the following, the Palais–Smale condition (cf. [10]) is referred to as (P-S). Now suppose that F satisfies the following conditions:

$$(S) \quad \inf_{x \in Z} F(x) > d > -\infty \text{ and } \lim_{\|y\| \rightarrow \infty} F(y) < d, \text{ for } y \in Y.$$

It then follows from (S) that

$$(I) \quad \text{there exists } r > 0 \text{ such that } \max_{y \in S(r)} F(y) < \inf_{z \in Z} F(z),$$

where $S(r) = \{y \in Y : \|y\| = r\}$. Let $\bar{\Gamma}$ be the family of sets $A \subset E$ satisfying the following properties:

(C1) For any continuous mapping $\sigma: A \rightarrow Y$ such that $\sigma(x) = x$ for all $x \in A \cap S$, the condition $0 \in \sigma(A)$ holds;

(C2) there exists a constant d such that for each integer $n \geq 1$, A can be covered by n^k balls of radius d/n .

We set

$$\bar{c} = \inf_{A \in \bar{\Gamma}} \max_{y \in A} F(y).$$

In [8], Lazer and Solimini established the following theorem, which we use to prove our main results.

THEOREM A. *Assume (I). If the (P-S) condition holds in \bar{c} and if F' is Fredholm of index 0 at all the critical points in $F^{-1}(\bar{c})$, then at least one of them has Morse index less than or equal to k and augmented Morse index greater than or equal to k .*

Remark 3. Let $z_0 \in Z$. Then the assertion of Theorem A holds if (S) and (I) are replaced by

$$(S') \quad \inf_{x \in Z} F(x) > d > -\infty \text{ and } \lim_{\|y\| \rightarrow \infty} F(y) < d, \text{ for } y \in z_0 + Y$$

and

$$(I') \quad \text{there exists } r > 0 \text{ such that } \max_{y \in S(r) + z_0} F(y) < \inf_{z \in Z} F(z).$$

In fact, if (S') and (I') are satisfied by F , the functional F_0 defined by $F_0(u) = F(u + z_0)$ satisfies (S) and (I).

3. PROOFS OF THE MAIN RESULTS

We put $\max G'' = \max\{|G''(x)| : x \in R^N\}$. Let $T_0 = 2/\sqrt{\max G''}$. In the following, we fix a positive number T such that $T < T_0$. Let $f \in \tilde{H}_T$. We put

$$S = \{u \in H_T : u \text{ is a solution of problem (P)}\}.$$

LEMMA 1. (1) S is bounded in H_T ;

(2) there exists $\rho > 0$ such that if $\|f\|_T \leq \rho$, then for each $u \in S$, there exists $i \in \{1, 2, \dots, m\}$ satisfying

$$|G''(u(t)) - G''(z_i)| < \frac{c_0}{2(N-1)} \quad \text{and} \quad |\langle G''(u(t))w_{i,j}, w_{i,j} \rangle| \geq c_0$$

for all $t \in [0, T]$ and $1 \leq j \leq N$.

Proof. Let $u \in H_T$ be a solution of (P). We multiply (P) by \tilde{u} and integrate over $[0, T]$. Then noting that $f \in \tilde{H}_T$ and $G'(u) = G'(\tilde{u}) + G''(\theta)(\tilde{u})$ for some mapping $\theta : [0, T] \rightarrow R^N$, we have

$$\|u_t\|_T^2 = \langle G'(u) - f, \tilde{u} \rangle_T \leq (\max G'')\|\tilde{u}\|_T^2 + \|f\|_T\|\tilde{u}\|_T. \quad (3.1)$$

Then since $\|\tilde{u}\|_T \leq (T/2\pi)\|u_d\|_T \leq (T/2)\|u_d\|_T$, we have that

$$\left(1 - \frac{T^2}{4} \max G''\right) \|u_d\|_T^2 \leq \|f\|_T \|\tilde{u}\|_T \leq \frac{T}{2} \|u_d\|_T \|f\|_T,$$

and then

$$\|\tilde{u}\|_T \leq \frac{T}{2} \|u_d\|_T \leq C \|f\|_T, \quad (3.2)$$

where $C = 1/((4/T^2) - \max G'')$. We next multiply (P) by $G'(u)$. Then we have by (3.1) and (3.2) that

$$\begin{aligned} \|G'(u)\|_T^2 &\leq \langle f, G'(u) \rangle_T + \|G''(u)u_d\|_T \|u_d\|_T \\ &\leq ((\max G'')\|\tilde{u}\|_T \|f\|_T + (\max G'')\|u_d\|_T^2) \leq C_1 \|f\|^2, \end{aligned} \quad (3.3)$$

where C_1 is a constant depending only on C and $\max G''$. It follows from (3.1) and (3.2) that there exists $\varepsilon > 0$ such that

$$|u(t) - \bar{u}| \leq \varepsilon \quad \text{for all } t \in [0, T]. \quad (3.4)$$

Then we have that $|\bar{u}| < d(\varepsilon)$. In fact, if $|\bar{u}| \geq d(\varepsilon)$, then we have by (2.6) that there exists $1 \leq i \leq N$ such that

$$G'(u(t))_{(i)} \cdot G''(\bar{u})_{(i)} > 0 \quad \text{for all } t \in [0, T].$$

This implies that

$$0 \neq \int_0^T G'(u) = \int_0^T f dt = 0.$$

This is a contradiction. Thus we have seen that assertion (1) holds. We next prove (2). By (3.2) and (3.3), we can choose $\rho > 0$ so small that if $\|f\|_T \leq \rho$, each solution $u \in H_T$ of (P) satisfies

$$|u(t) - \bar{u}| \leq \varepsilon_0/2 \quad \text{for all } t \in [0, T] \quad (3.5)$$

and

$$|G'(u(t))| \leq m_0/2 \quad \text{for all } t \in [0, T]. \quad (3.6)$$

Now suppose that $\|f\|_T \leq \rho$. From the argument above, we have that $|\bar{u}| \leq d(\varepsilon_0)$. If $\bar{u} \notin \bigcup (z_i + D(\varepsilon_0/2))$, then by (2.8) and (3.5), we find that $\sup_{t \in [0, T]} |G'(u(t))| \geq m_0$. This contradicts (3.6). Thus we have seen that $\bar{u} \in \bigcup (z_i + D(\varepsilon_0/2))$. Then assertion (2) follows from (3.5) and (2.7). ■

Throughout the rest of this section, we assume that $\|f\|_T \leq \rho$. For each positive integer k , we define a functional $F_k: H_{kT} \rightarrow \mathbb{R}$ by

$$F_k(u) = \int_0^{kT} \left(\frac{1}{2} |u_t|^2 - G(u) + \langle f(t), u(t) \rangle \right) dt \quad \text{for } u \in H_{kT}.$$

It is then easy to see that each critical point of F_{kT} is a kT -periodic solution of (P). We denote by $\text{m-index}_k(u)$ ($\text{am-index}_k(u)$) the Morse index (augmented Morse index) of F_k at a critical point $u \in H_{kT}$ of F_k . Let Γ_k be the set $\bar{\Gamma}$ defined in Section 2 with E , Y , and Z replaced by H_{kT} , \bar{H}_{kT} , and \tilde{H}_{kT} , respectively. We set

$$c_k = \inf_{A \in \Gamma_k} \max_{y \in A} F_k(y).$$

LEMMA 2. For each $k \geq 1$, F_k satisfies (I') with E , Y , and Z replaced by H_{kT} , \bar{H}_{kT} , and \tilde{H}_{kT} .

Proof. Let $k \geq 1$. Let $u_0 \in \tilde{H}_T$ be the unique element satisfying

$$m_1 = \langle f, u_0 \rangle_T + \frac{1}{2} \|u_0\|_T^2 = \min \left\{ \langle f, v \rangle_T + \frac{1}{2} \|v\|_T^2 : v \in \tilde{H}_T \right\}.$$

Then since $f \in \tilde{H}_T$, we find that

$$m_1 k = \min \left\{ \langle f, v \rangle_{kT} + \frac{1}{2} \|v\|_{kT}^2 : v \in \tilde{H}_{kT} \right\}.$$

For each $r > 0$, we set

$$S_0(r) = \{x + u_0 : x \in \mathbb{R}^N, |x| = r\}.$$

Then it follows from (G1) and the definition of u_0 that

$$\limsup_{r \rightarrow \infty} \{F_k(v) : v \in u_0 + \mathbb{R}^N \setminus D(r)\} = M_k = - \left(\lim_{|x| \rightarrow \infty} G(x) \right) kT + m_1 k. \quad (3.7)$$

Here let C be a positive number such that

$$\|\tilde{v}\|_{kT} \leq C\|v_t\|_{kT} \quad \text{for } v \in H_{kT}.$$

By (G1), we can choose $c_1, c_2 > 0$ such that $c_1 C^2 < 1/2$ and

$$|G(x)| \leq c_1|x|^2 + c_2 \quad \text{for all } x \in \mathbb{R}^N.$$

Then we have by (G1) and the inequalities above that

$$\begin{aligned} F_k(v) &= \int_0^{kT} \left(\frac{1}{2} \|v_t\|^2 - G(v) + \langle f, v \rangle \right) dt \\ &> \left(\left(\frac{1}{2} - c_1 C^2 \right) \|v_t\|_{kT}^2 - C\|f\|_{kT}\|v_t\|_{kT} - c_2 kT \right) \end{aligned}$$

for $v \in \tilde{H}_{kT}$. Then there exists $d > 0$ such that

$$\inf\{F_k(v) : v \in \tilde{H}_{kT}, \|v_t\| \geq d\} > M_k. \quad (3.8)$$

On the other hand, noting that $\sup\{|v(t)| : v \in \tilde{H}_{kT}, \|v_t\| \leq d\} < \infty$, we can see that

$$\sup \left\{ \int_0^{kT} G(v) : v \in \tilde{H}_{kT}, \|v_t\| \leq d \right\} = M_d < \left(\lim_{|x| \rightarrow \infty} G(x) \right) kT.$$

Then we have

$$F_k(v) = \int_0^{kT} \left(\frac{1}{2} \|v_t\|^2 - G(v) + \langle f, v \rangle \right) dt > -M_d - m_1 k \quad (3.9)$$

for all $v \in \tilde{H}_{kT}$ with $\|v_t\| \leq d$. Consequently, we have by (3.8) and (3.9) that

$$\inf\{F_k(v) : v \in \tilde{H}_{kT}\} = \tilde{M}_k > M_k. \quad (3.10)$$

Therefore F_k satisfies (S') with E , Y , and Z replaced by H_{kT} , \overline{H}_{kT} , and \tilde{H}_{kT} and $z_0 = u_0$. Then for each $k \geq 1$, there exists $r_k > 0$ such that $\sup\{F_k(v) : v \in S_0(r_k)\} < \tilde{M}_k$. That is F_k satisfies (I') with $S(r)$ replaced by $S_0(r_k)$. ■

LEMMA 3. For each $k \geq 1$, the following assertions hold.

- (i) $c_k > M_k$;
- (ii) F_k satisfies (P-S) at c_k .

Proof. Let $k \geq 1$. (i) Let P be the projection from H_{kT} onto \overline{H}_{kT} . Then from the definition of Γ_k , $0 \in P(A)$ for any $A \in \Gamma_k$. This implies that $A \cap \tilde{H}_{kT} \neq \emptyset$ for any $A \in \Gamma_k$. Then (i) follows from (3.10). (ii) Let $\{u_n\} \in H_{kT}$ be a sequence such that $F'_k(u_n) \rightarrow 0$ and $F_k(u_n) \rightarrow c_k$, as $n \rightarrow \infty$. Let C be the positive constant defined in the proof of Lemma 2. Then we have

$$\begin{aligned} F_k(u_n) &= \int_0^{kT} \left(\frac{1}{2} \|\tilde{u}_{nt}\|^2 - G(u_n) + \langle f(t), \tilde{u}_n(t) \rangle \right) dt \\ &\geq (1/2C) \|\tilde{u}_n\|_{kT}^2 - \int_0^{kT} G(u_n) dt - \|f\|_{kT} \|\tilde{u}_n\|_{kT}. \end{aligned} \quad (3.11)$$

Suppose that $\limsup_n \|\tilde{u}_n\|_{kT} = \infty$. If there exists $C_1 > 0$ and

$$\limsup_n \|\tilde{u}_n\|_{kT} / \|\tilde{u}\|_{kT} = C_1 < \infty,$$

then nothing that $\|u_n\|_{kT} \leq (2C_1 + 1)\|\tilde{u}_n\|$ for n sufficiently large, and $\lim_{|x| \rightarrow \infty} G(x)/|x|^2 = 0$, we have from (2.11) that $\limsup F_k(u_n) = \infty$. This is a contradiction. Therefore we have that $\limsup_n \|\tilde{u}_n\|_{kT} / \|\tilde{u}_n\|_{kT} = \infty$. From the assumption, we have by applying $F'_k(u_n)$ to $\tilde{u}_n / \|\tilde{u}_n\|_{kT}$ that

$$\lim_{n \rightarrow \infty} \left(\|\tilde{u}_{nt}\|_{kT}^2 / \|\tilde{u}_n\|_{kT} - \int_0^{kT} \langle G'(u_n) - f, \tilde{u}_n / \|\tilde{u}_n\|_{kT} \rangle dt \right) = 0. \quad (3.12)$$

On the other hand, for each $n \geq 1$, there exists a mapping $\theta_n: [0, kT] \rightarrow R^N$ such that

$$\theta_n(t) \in \{\bar{u}_m(t) + s\tilde{u}_n(t) : s \in [0, 1]\} \quad \text{for each } t \in [0, kT]$$

and

$$G'(u_n) = G'(\bar{u}_n) + G''(\theta_n)\tilde{u}_m \quad \text{for each } n \geq 1.$$

Then since \tilde{u}_n has mean value zero, we have that

$$\int_0^{kT} \langle G'(u_n), \tilde{u}_n \rangle dt = \int_0^{kT} \langle G''(\theta_n)\tilde{u}_n, \tilde{u}_n \rangle dt. \quad (3.13)$$

Since $\lim_{n \rightarrow \infty} \|\tilde{u}_n\|_{kT}/\|\tilde{u}_n\|_{kT} = \infty$, we have that $\lim_{n \rightarrow \infty} |\theta_n(t)|_{kT} = \infty$ on $[0, kT]$ and then $\lim_{n \rightarrow \infty} G'(\theta_n(t)) = 0$ for all $t \in [0, kT]$. Therefore recalling that $\|\tilde{u}_n\|_{kT} \leq C\|\tilde{u}_{nt}\|$, we have by (3.13) that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left(\|\tilde{u}_{nt}\|_{kT}^2 / \|\tilde{u}_n\| - \int_0^{kT} \langle G'(u_n) - f, \tilde{u}_n / \|\tilde{u}_n\|_{kT} \rangle dt \right) \\ & \geq \lim_{n \rightarrow \infty} \left((1/C^2) \|\tilde{u}_n\|_{kT} - \int_0^{kT} \langle G'(\theta_n) - f, \tilde{u}_n / \|\tilde{u}_n\|_{kT} \rangle dt \right) \\ & \geq \lim_{n \rightarrow \infty} \left((1/C^2) \|\tilde{u}_n\|_{kT} - \int_0^{kT} \langle G'(\theta_n), \tilde{u}_n / \|\tilde{u}_n\|_{kT} \rangle dt - \|f\|_{kT} \right). \end{aligned}$$

This contradicts (3.12). Consequently, we obtain that $\{\tilde{u}_n\}$ is bounded in H_{kT} . We next see that $\{\tilde{u}_n\}$ is bounded. Suppose that $\{\tilde{u}_n\}$ is unbounded. We may assume by extracting a subsequence that $\lim \|\tilde{u}_n\| = \infty$. It then follows that

$$\lim_{n \rightarrow \infty} \|G'(u_n)\|_{kT} = 0. \quad (3.14)$$

In fact, if $\limsup \|G'(u_n)\| > 0$, then recalling that $\{\tilde{u}_n\}$ is bounded, we have by (2.6) that

$$\limsup_n \left| \int_0^{kT} G'(u_n) dt \right| > 0.$$

Since

$$\lim_{n \rightarrow \infty} F'_k(u_n)(-1) = \lim_{n \rightarrow \infty} \int_0^{kT} (G'(u_n) - f) dt = \lim_{n \rightarrow \infty} \int_0^{kT} G'(u_n) dt = 0,$$

this is a contradiction. Here we define a convex functional F_0 by

$$F_0(v) = \int_0^{kT} \left(\frac{1}{2} \|v_t\|^2 + \langle f, v \rangle \right) dt \quad \text{for } v \in H_{kT}.$$

Then we have that

$$\begin{aligned} F'_k(u_n)(v) &= \int_0^{kT} (\langle u_{nt}, v_t \rangle - \langle G'(u_n), v \rangle + \langle f, v \rangle) dt \\ &= F'_0(u_n)(v) - \int_0^{kT} \langle G'(u_n), v \rangle dt \quad \text{for } v \in H_{kT}. \end{aligned}$$

Recalling that $F'_k(u_n) \rightarrow 0$ as $n \rightarrow \infty$, we find by (3.14) and the equality above that $F'_0(u_n) \rightarrow 0$ as $n \rightarrow \infty$. It is then easy to see from the definition of functional F_0 that $\{u_n\}$ is a minimizing sequence of the functional F_0 . This implies that

$$\lim_{n \rightarrow \infty} \left(\langle f, u_n \rangle_{kT} + \frac{1}{2} \|u_n\|_{kT}^2 \right) = k \cdot m_1.$$

Therefore from (G1), we obtain that

$$\lim_{n \rightarrow \infty} F_k(u_n) = \lim_{n \rightarrow \infty} \left(\frac{1}{2} \|u_n\|_{kT}^2 + \langle f, u_n \rangle_{kT} - \left(\lim_{|x| \rightarrow \infty} G(x) \right) kT \right) = M_k < c_k.$$

This contradicts the assumption. Thus we obtain that $\{\bar{u}_n\}$ is bounded. Then we can see by the standard argument (cf. [10]) that $\{u_n\}$ has a convergent subsequence. This completes the proof. ■

Remark 4. The nondegeneracy of G is used only for proving (2) of Lemma 1. Then (1) of Lemma 1, Lemma 2, and Lemma 3 are valid without the nondegeneracy.

Let $u \in S$. Then by (2) of Lemma 1, there exists $1 \leq i \leq m$ such that

$$|\langle G''(u(t))w_{i,j}, w_{i,j} \rangle| \geq c_0 \quad \text{for all } t \in [0, T] \text{ and } 1 \leq j \leq N.$$

Then since $G \in C^2(R^N, R)$ and $u: [0, T] \rightarrow R^N$ is continuous, this implies that for each $u \in S$, there exists $1 \leq i \leq m$ such that for each $1 \leq j \leq N$,

$$\langle G''(u(t))w_{i,j}, w_{i,j} \rangle \geq c_0 \quad \text{on } [0, T]$$

or

$$\langle G''(u(t))w_{i,j}, w_{i,j} \rangle \leq -c_0 \quad \text{on } [0, T]$$

holds. Here we put

$$\bar{\mu}_i(u) = \max_{1 \leq j \leq N} \sup \{ \langle G''(u(t))w_{i,j}, w_{i,j} \rangle, t \in [0, T] \}$$

and

$$\underline{\mu}_i(u) = \max_{1 \leq j \leq N} \inf\{\langle G''(u(t))w_{i,j}, w_{i,j} \rangle, t \in [0, T]\}$$

for each $1 \leq i \leq m$. We also set

$$S_1 = \{u \in S : \underline{\mu}_i(u) \geq c_0 \text{ for some } 1 \leq i \leq m\}$$

$$S_2 = \{u \in S : \bar{\mu}_i(u) \leq -c_0 \text{ for some } 1 \leq i \leq m\}.$$

Then it follows from the observation above that $S = S_1 \cup S_2$.

PROPOSITION 1. *There exists a positive integer k_0 such that for each $k \geq k_0$,*

$$m\text{-index}_k(u) > N \quad \text{for all } u \in S_1.$$

Proof. Let $u \in S_1$. Then from the definition of S_1 , we have that there exist $1 \leq i \leq m$ and $1 \leq j \leq N$ such that

$$\langle G''(u(t))w_{i,j}, w_{i,j} \rangle \geq c_0 \quad \text{for all } t \in [0, T].$$

Here we put $w = w_{i,j}$. Then from the inequality above

$$(1/T) \int_0^T \langle G''(u(t))w, w \rangle dt \geq c_0.$$

Here we put $d_{mn} = (1/T) \int_0^T (G''(u(t)))_{mn} dt$ for each $1 \leq m, n \leq N$, where $(G''(u(t)))_{mn}$ is the (m, n) -element of the matrix $G''(u(t))$. Then from the inequality above, we find that the matrix $D = \{d_{mn}\}$ satisfies

$$\langle Dw, w \rangle \geq c_0. \quad (3.15)$$

We note that D is Hermitian. We define a matrix $A = \{a_{mn}\}$ by $a_{mn}(t) = (G''(u(t)))_{mn} - d_{mn}$ for $m, n \in \{1, \dots, N\}$. Then each a_{mn} is a T -periodic function with $\bar{a}_{mn} = 0$. From the definitions of D and A , we have

$$G''(u) = D + A.$$

By (3.15), D has a positive eigenvalue $\lambda (\geq c_0)$. Let $y = (y_1, \dots, y_N) \in \mathbb{R}^N$ be a normalized eigenvector corresponding to the eigenvalue λ . Here we choose a positive prime number k_0 such that

$$(2\pi(N+1)/k_0T)^2 < \lambda \quad \text{and} \quad 2(N+1) < k_0. \quad (3.16)$$

Now let $k \geq k_0$, and

$$v_i(t) = \left(\sqrt{\frac{2}{kT}} \sin\left(\frac{2\pi i}{kT}t\right) \right) y \quad \text{for } 1 \leq i \leq N+1.$$

Then since each a_{mn} is T -periodic and $\bar{a}_{mn} = 0$, we have that each n th element of Av_i consists of the linear combination of functions $\sin(2\pi(kj \pm i)t/kT)$, $\cos(2\pi(kj \pm i)t/kT)$, $j = 1, 2, \dots$. From the second inequality in (3.16), we find that these functions are orthogonal to $\sin((2\pi qt/kT))$, $1 \leq q \leq N+1$ in $L^2(0, kT)$. It then follows that

$$\langle Av_i, v_j \rangle_{kT} = 0 \quad \text{for all } 1 \leq i, j \leq N+1.$$

Then we find that

$$\begin{aligned} \langle -v_{iii} - G''(u)v_i, v_i \rangle_{kT} &= (2\pi i/kT)^2 - \langle Dv_i, v_i \rangle_{kT} \\ &= (2\pi i/kT)^2 - \langle Dy, y \rangle \\ &= (2\pi i/kT)^2 - \lambda < 0 \end{aligned} \quad (3.17)$$

for all $1 \leq i \leq N+1$. On the other hand, recalling that $2(N+1) < k$ and each a_{mn} is T -periodic, we can see that

$$\langle G''(u)v_i, v_j \rangle_{kT} = \langle (A + D)v_i, v_j \rangle_{kT} = 0 \quad \text{for } i \neq j.$$

Then it easy to see from (3.17) and the equality above that for any $v \in \text{span}\{v_1, \dots, v_{N+1}\} \setminus \{0\}$,

$$\langle -v_{iii} - G''(u)v, v \rangle_{kT} < 0.$$

This implies that $\text{m-index}_k(u) > N$ for $k \geq k_0$. ■

PROPOSITION 2. For each $k \geq 1$,

$$\text{am-index}_k(u) = 0 \quad \text{for all } u \in S_2.$$

Proof. Let $u \in S_2$. That is, there exists an orthonormal basis $\{w_{i,1}, w_{i,2}, \dots, w_{i,N}\}$ of R^N such that

$$\langle G''(u(t))w_{i,j}, w_{i,j} \rangle \leq c_0 \quad \text{on } [0, T] \text{ for all } 1 \leq j \leq N. \quad (3.18)$$

Since $|G''(u(t)) - G''(z_i)| < c_0/2(N-1)$ on $[0, T]$ by the first inequality of (2) of Lemma 1, we find that for each $j, k \in \{1, \dots, N\}$ with $i \neq j$,

$$\begin{aligned}
& |\langle G''(u(t))w_{i,j}, w_{i,k} \rangle| \\
& \leq |\langle (G''(u(t)) - G''(z_i))w_{i,j}, w_{i,k} \rangle + \langle G''(z_i)w_{i,j}, w_{i,k} \rangle| \quad (3.19) \\
& \leq \|G''(u(t)) - G''(z_i)\|w_{i,j}\|w_{i,k}\| \leq c_0/2(N-1).
\end{aligned}$$

Then it follows from (3.18) and (3.19) that

$$\langle G''(u(t))w, w \rangle \leq -c_0/2 \quad \text{for all } w \in R^N \text{ with } |w| = 1 \text{ and } t \in [0, T].$$

Then we have that

$$\langle -v_{tt} - G''(u)v, v \rangle_{kT} \geq \|v_t\|_{kT}^2 + (c_0/2)\|v\|_{kT}^2 \quad \text{for all } v \in H_{kT}.$$

Then the assertion follows. ■

Now we can complete the proof of Theorem 1.

Proof of Theorem 1. Let k be a prime number with $k \geq k_0$. Then by applying Theorem A to the functional F_k , we find that there exists a critical point u_k of F_k satisfying that

$$\text{am-index}_k(u) \geq N \quad \text{and} \quad \text{m-index}_k(u) \leq N.$$

Since k is a prime number, the minimal period of u_k is T or kT . If the minimal period of u_k is T , we have that $u_k \in S = S_1 \cup S_2$. Then by Proposition 1 and Proposition 2, we find that $\text{m-index}_k(u) > N$ or $\text{am-index}_k(u) = 0$ holds. This is a contradiction. Therefore the minimal period of u_k is kT . This completes the proof. ■

Proof of Theorem 2. By Remark 4, we have that S is bounded in H_T by (1) of Lemma 1. Then we have by (G3) that there exists $d > 0$ such that

$$\langle G''(u(t))x_0, x_0 \rangle \geq d \quad \text{for all } u \in S \text{ and } t \in [0, T].$$

Then the argument of Proposition 1 is valid for all $u \in S$ with $w \in R^N$ and c_0 replaced by x_0 and d , respectively. That is, we have that there exists a positive integer such that for each $k \geq k_0$, $\text{m-index}_k(u) > N$ for all $u \in S$. Then the assertion of Theorem 2 follows from the argument of the proof of Theorem 1. ■

Proof of Theorem 3. The restriction that $T < T_0$ is used only in inequality (3.2) to obtain the boundedness of $\tilde{S} = \{\tilde{u} : u \in S\}$. If G' is bounded, then \tilde{S} is bounded for any $T > 0$ (see the proof of Theorem 1 of [5]). Then the assertion follows by the argument for Theorem 1 and Theorem 2. ■

ACKNOWLEDGMENT

The author wishes to express his thanks to the referee for reading this article very carefully and giving useful comments.

REFERENCES

1. E. ALVES DE B. E SILVA, "Critical Point Theorems and Applications to Differential Equations," Ph.D. thesis, University of Wisconsin, Madison, 1988.
2. F. CLARKE AND I. EKELAND, Nonlinear oscillations and boundary value problems for Hamiltonian systems, *Arch. Rational Mech. Anal.* **78** (1982), 315–333.
3. I. EKELAND AND H. HOFER, Subharmonics for convex nonautonomous Hamiltonian systems, *Comm. Pure Appl. Math.* **40** (1987), 1–36.
4. A. FONDA AND M. WILLEM, Subharmonic oscillations of forced pendulum-type equations, *J. Differential Equations* **81** (1989), 215–220.
5. A. FONDA AND A. C. LAZER, Subharmonic solutions of conservative systems with nonconvex potentials, *Proc. Amer. Math. Soc.* **115** (1992), 183–190.
6. A. FONDA, M. RAMOS, AND M. WILLEM, Subharmonic solutions for second order differential equations, *Top. Meth. Non. Anal.* **1** (1993), 49–66.
7. F. GIANNONI, Periodic solutions of dynamical systems by the saddle point theorem of Paul H. Rabinowitz, *Nonlinear Anal.* **12** (1988), 761–775.
8. A. C. LAZER AND S. SOLIMINI, Nontrivial solutions of operator equations and Morse indices of critical points of min–max type, *Nonlinear Anal.* **12** (1988), 761–775.
9. J. MAWHIN AND M. WILLEM, "Critical Point Theory and Hamiltonian Systems," Springer-Verlag, New York, 1988.
10. P. H. RABINOWITZ, "Minimax Methods in Critical Point Theory with Applications to Differential Equations," CBMS Reg. Conf. Ser. in Math. No 65, Amer. Math. Soc., Providence, RI, 1986.
11. P. H. RABINOWITZ, On subharmonic solutions of Hamiltonian systems, *Comm. Pure Appl. Math.* **33** (1990), 608–633.
12. S. SOLIMINI AND S. TERRACINI, A multiplicity result for periodic solutions of autonomous conservative systems, *Nonlinear Anal.* **16** (1992), 903–916.
13. M. WILLEM, Subharmonic oscillations of convex Hamiltonian systems, *Nonlinear Anal.* **9** (1985), 1303–1311.